



Medical Policy

Subject:	Voretigene neparvovec-rzyl (Luxturna®)	Publish Date:	04/24/2019
Document#:	MED.00120	Last Review Date:	03/21/2019
Status:	Reviewed		

Description/Scope

This document addresses the use of voretigene neparvovec-rzyl (Luxturna, Spark Therapeutics, Philadelphia, PA), a gene replacement therapy intended to treat retinal dystrophies caused by biallelic RPE65 gene mutations.

Note: Please see the following related document regarding genetic testing for individuals with inherited diseases, including inherited retinal diseases:

- CG-GENE-13 Genetic Testing for Inherited Diseases

Position Statement

Medically Necessary:

The use of voretigene neparvovec-rzyl gene therapy is considered **medically necessary** in individuals who meet *all* of the following criteria:

- A. A diagnosis of retinal dystrophy due to confirmed *RPE65* mutation(s) in *both* alleles; **and**
 - B. At least 1 year of age; **and**
 - C. For each eye indicated for treatment, sufficient viable retinal cells as determined by non-invasive means, such as optical coherence tomography (OCT) and/or ophthalmoscopy as evidenced by *one* of the following:
 1. An area of retina within the posterior pole of greater than 100 µm thickness shown on OCT; **or**
 2. Three or more disc areas of retina without atrophy or pigmentary degeneration within the posterior pole; **or**
 3. Remaining visual field within 30 degrees of fixation as measured by a III4e isopter or equivalent;
- and**
- D. For each eye indicated for treatment, have not had intraocular surgery within 6 months.

Note: Voretigene neparvovec-rzyl has not been adequately studied in pregnant women or individuals unwilling to use contraception for four months after treatment.

Investigational and Not Medically Necessary:

The use of voretigene neparvovec-rzyl gene therapy is considered **investigational and not medically necessary** when the above criteria are not met.

Repeat injections of voretigene neparvovec-rzyl gene therapy in the same eye are considered **investigational and not medically necessary** in all cases.

Rationale

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Voretigene neparvovec-rzyl (Luxturna)

On December 19, 2017 the U.S. Food and Drug Administration (FDA) approved voretigene neparvovec-rzyl (Luxturna), the first gene replacement (MAF) therapy approved in the U.S. to treat an inherited disease. Voretigene neparvovec-rzyl is an adeno-associated virus vector-based gene therapy for the treatment of biallelic RPE65 mutation-associated retinal dystrophy. Voretigene neparvovec-rzyl is supplied in single dose vials to be used in a single eye and is administered via subretinal injection. Treatment must be done separately in each eye on separate days, with at least six days between procedures. Use of this therapy is limited to those with evidence of viable retinal cells and to those age 12 months or older. Infants younger than 12 months of age are undergoing active retinal cell proliferation and there is a potential for dilution or loss of voretigene neparvovec-rzyl after administration in this population.

FDA approval of voretigene neparvovec-rzyl was primarily based on the results of an open-label, randomised, controlled phase 3 trial (Russell, 2017). A total of 29 individuals age 3 years or older with Leber congenital amaurosis type 2 (LCA2) due to confirmed biallelic *RPE65* gene mutations with both eyes with a visual acuity of 20/60 or worse or visual field less than 20 degrees in any meridian or both, and sufficient viable retinal cells present in each eye participated in the study. The intervention group received a subretinal injection of voretigene neparvovec-rzyl in the poorer seeing eye, followed by an injection into the second eye 6-18 days after the initial injection. A control group of 9 individuals did not receive therapy for 1 year at which time they could cross over and receive recombinant adeno-associated virus voretigene neparvovec-rzyl therapy (AAV2-hRPE65v2). The primary efficacy endpoint was the change from baseline to 1 year in the score of the multi-luminance mobility testing (MLMT), an assessment of functional vision at specified light levels. Secondary efficacy endpoints also included full-field light sensitivity threshold (FST) testing and best-corrected visual acuity (BCVA), each averaged over both eyes. At 1 year, the mean bilateral MLMT score improvement in the intervention group was 1.8 (standard deviation [SD] 1.1) compared to 0.2 (SD 1.0) in the control group for a difference of 1.6 (95% confidence interval [CI], 0.72-2.41, p=0.0013). Mean FST scores in the intervention group improved by 30 days following intervention and remained stable at 1 year. There was no meaningful change in FST scores in the control group at 1 year. The BCVA scores in the intervention group did show an insignificant numerical improvement over the control group. The authors noted this was not unexpected as BCVA is a measure of foveal, cone-mediated function, and the *RPE65* gene mutations result in a rod-mediated disease. The most common treatment-related ocular adverse events included elevated intraocular pressure, cataract and eye inflammation and were considered mild or moderate. There were no reported serious product-related adverse events or harmful immune responses. One individual in the intervention group did experience loss of visual acuity in the first eye which was significant based upon the VA measurement criteria, however, the individual did not report a change in vision. The authors noted the loss might be due to foveal thinning following subretinal injection, however, the analysis was inconclusive.

In an open label phase 1-2 trial, the safety and efficacy of gene therapy with a recombinant adeno-associated virus 2/2 (rAAV2/2) vector carrying the *RPE65* complementary DNA was evaluated for 3 years following treatment (Bainbridge, 2015). A total of 12 children and young adults (6 to 23 years old) with early-onset, severe retinal dystrophy, caused by mutations in *RPE65*, were treated with RPE65 gene supplementation. Individuals received treatment in 1 eye; 4 individuals received a lower dose while the remaining 8 individuals received a higher dose. In addition to a baseline evaluation, individuals underwent interval evaluations for 3 years, measuring visual acuity, contrast sensitivity, color vision, and spectral sensitivities. While there was a greater proportion of individuals with improved retinal sensitivity among those treated with higher versus lower doses, due to the widely differing level of improvement among the individuals and the small number of participants, a dose-response effect may be present but not definitively concluded. Additionally, treatment benefits did not appear to be lasting. Retinal sensitivity improvement was reported within 1 to 2 months following therapy with progressive improvement

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Medical Policy

Voretigene neparvovec-rzyl (Luxturna)

continuing for 6 to 12 months. Following 12 months, retinal sensitivity declined in the majority of cases, although 2 individuals showed a continued improvement over pre-intervention levels over 3 years. Visual acuity was improved in 1 individual who showed a similar improvement in the contralateral untreated eye. The authors noted rAAV2/2 vector carrying the *RPE65* cDNA gene therapy led to a “temporary, variable and incomplete restoration of retinal function in humans.”

Bennett and colleagues (2016) assessed the safety and efficacy of administering the treatments in the contralateral eye in a follow-up to the initial study. A total of 11 of the 12 original participants received an injection in the contralateral, untreated eye and were evaluated on a regular basis for 3 years. An individual with glaucoma in the uninjected eye could not be included in this follow-up study. One individual developed bacterial endophthalmitis following the injection. The remaining adverse effects related to the procedure included dellen formation (n=2) and cataracts (n=1), all of which were successfully treated. For the majority of outcomes, results for the second injected eye were compared with results over the same follow-up period for the initial injected eye. In a pooled analysis, results from full-field light sensitivity threshold testing showed robust improvements in both rod and cone function in the contralateral eyes by day 30 through year 3. A pooled analysis also did not show any significant changes in visual acuity from baseline through year 3 in either eye. There were reported significant improvements in mobility testing and pupillary light reflex testing results which lasted through year 3.

Weleber and associates (2016) treated 12 individuals in a small, non-randomized multi-center clinical trial. A total of 8 adults and 4 children with either LCA or severe early-childhood-onset retinal degeneration (SECORD) with varying degrees of visual acuity were included in the study. Individuals received a subretinal injection containing either 1.8×10^{11} (group 1) or 6×10^{11} (group 2) vector genomes of rAAV2-CBhRPE65. While 9 of the 12 individuals showed an improvement in at least 1 measure of visual function, the specific area and amount of improvement varied among individuals. Two individuals reported a decrease in BCVA or visual field area that was greater in the treated eye compared to the untreated eye. Adverse events related to the study agent included ocular hyperemia (n=2) and photopsia (n=1). The authors noted that larger studies with longer follow-up is needed to further assess side effects or safety concerns.

Le Meur and colleagues (2018) evaluated the efficacy and tolerance of injections of AAV serotypes 2 and 4 (AAV2/4) RPE65 in an RPE65 gene deficiency LCA in a phase 1/2 study. A total of 9 individuals age 9 to 42 years with 2 confirmed RPE65 gene mutations were grouped into 3 cohorts based upon age and injected viral vector dose. Cohort 1 consisted of adults who received the lowest dose of viral vector, cohorts 2 and 3 received higher doses of viral vector. Each participant received 2 to 5 simultaneous unilateral subretinal injections with dosages between 1.22×10^{10} to 4.8×10^{10} vector genomes (vg) into the eye with the greatest visual impairment. All 9 individuals were followed for 1 year and 6 of the individuals agreed to additional follow-up for 2 to 3.5 years. Visual acuity loss progression varied among the cohorts. In cohorts 1 and 3, there were no observed alterations in visual acuity. Cohort 2 showed a trend towards improved visual acuity. Visual acuity in the treated eye remained stable during the follow-up period. Visual acuity in the untreated eye trended toward reduced visual acuity. However, none of the alterations attained statistical significance. Individuals who presented with nystagmus and intermediate initial visual acuity reported the most marked improvements (cohort 2), while individuals in cohort 3 with correct initial visual acuity without nystagmus showed less of an improvement. The authors postulated that this may be related to the clinical modification in nystagmus. Changes in mobility, evaluated using travel time through a maze under different lighting conditions, were not significantly changed from pre-injection through 1 year follow-up. The injections were well tolerated with only transient symptoms reported. An analysis of the mean retinal thickness at 1, 2, or 3

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Voretigene neparvovec-rzyl (Luxturna)

year follow-up revealed no alterations in retinal thickness in the treated or untreated eyes. The authors noted that the possibility of retinal thinning may occur and anticipate assessing this factor in further follow-up.

Le Meur and associates compared their current study to previous studies. The researchers increased the number of injection subretinal sites (2-5) compared to earlier studies in an attempt to increase the contact between the viral vector and the retinal surface, but it remains unclear whether this approach provided additional benefit. In addition, the dose increments in this study were smaller than those in previous trials (1.22×10^{10} to 4.8×10^{10} versus 1.5×10^{10} vg to 6.11×10^{11} vg, respectively). The authors noted that the dose used may not have been sufficient to achieve a meaningful response or to prevent further deterioration. The authors concluded that long term follow-up is needed to evaluate the persistence of the effects and evaluate the treatment effect on retinal degeneration.

Treatment with voretigene neparvovec-rzyl is limited to individuals at least 1 year old. The Luxturna package insert (PI) label includes information regarding use in the pediatric population:

Treatment with LUXTURNA is not recommended for patients younger than 12 months of age, because the retinal cells are still undergoing cell proliferation, and LUXTURNA would potentially be diluted or lost during cell proliferation.

Early studies have shown safety and variable efficacy with variable duration for up to 3 years following injection therapy (Cideciyan, 2008; Le Meur, 2018; Jacobson, 2012; Maguire, 2009; Testa, 2013). There appears to be a limited or no systemic immune response to the subretinal AAV2 vector delivery (Jacobson, 2012). There remain several unknowns related to gene therapy, including which individuals would benefit from therapy and the most efficacious delivery method. In addition, ongoing durability remains a concern (Bennett, 2016; Le Meur, 2017; Webeler, 2016). At the current time, the safety and efficacy of repeat injections into the same eye have not been evaluated in clinical studies.

Background/Overview

Hereditary retinal dystrophies or inherited retinal diseases (IRDs) are progressive disorders which typically result in blindness at a young age. There are approximately 220 mutations associated with IRDs including the *RPE65* mutation (Russell, 2017). Missense mutations in the retinal pigment epithelium-specific protein 65 kDa (RPE65) result in a more severe form of the retinal degenerative disease that is associated with photoreceptor dysfunction and degeneration (Cideciyan, 2013). A lack of functional RPE65 results in rod photoreceptor cells that are unable to respond to light, contributing to decreased ability to see in low light conditions and diminished visual fields. As retinal degeneration progresses and affects cones, the disorder leads to blindness (FDA, 2017).

IRDs attributed to biallelic RPE65 mutations are associated with the subtypes of LCA type 2 and retinitis pigmentosa (RP) type 20. In the United States (U.S.) there are an estimated 1000 to 3000 individuals with biallelic RPE65 mutation-associated retinal dystrophy (FDA, 2017). Prior to the development of gene therapy, there were no pharmacologic treatments available to treat IRDs.

The 2016 American Academy of Ophthalmology (AAO) recommends genetic testing be ordered at the initial visit for individuals with a suspected inherited retinal degenerative disease. The causative mutation can be identified in up to 60-80% of affected individuals, which can guide treatment decisions. The scope of genetic testing recommended varies, multi-gene testing may be necessary when there are multiple causative genes, while single

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Voretigene neparvovec-rzyl (Luxturna)

gene analysis might be more appropriate for certain conditions. For diseases such as LCA, which is caused by multiple different genes, it can be more efficient to order a single test which has been designed to specifically evaluate for all of the known causative genes (Stone, 2012). A listing of the genes causing retinal disease can be accessed here: <https://sph.uth.edu/retnet/>.

Individuals with LCA caused by *RPE65* retain a largely normal outer retinal structure as compared to the amount of visual loss, a unique characteristic which makes potentially restoring vision via gene replacement therapy viable (Wright, 2015). However, gene replacement therapy has been shown to provide a partial improvement in visual dysfunction, as evidenced by improved vision, but does not appear to slow photoreceptor degeneration.

Treatment with voretigene neparvovec-rzyl involves the use of a standard three-port pars plana vitrectomy with removal of the posterior cortical vitreous followed by an injection of the vector genome of adeno-associated virus 2 (AAV2).hRPE65v2 into the subretinal space, creating a localized dome-shaped retinal detachment. The localized retinal detachment typically resolves within 14 to 24 hours following surgery. Minimal persistent subretinal fluid at the macula typically resolves within 2 to 3 days.

Warnings and Precautions

Warnings from the FDA PI Label (2017) include the following:

- Endophthalmitis: Use proper aseptic injection technique and monitor for signs and symptoms of infection.
- Permanent decline in visual acuity: Monitor for visual disturbances.
- Retinal abnormalities: Monitor for macular abnormalities, retinal tears or breaks. Do not inject in the immediate vicinity of the fovea.
- Increased intraocular pressure: Monitor and manage intraocular pressure elevations.
- Expansion of intraocular air bubbles: Air travel and/or scuba diving is not recommended until any intraocular air bubbles have been absorbed.
- Cataract: Subretinal injection of Luxturna may result in cataract formation or increase in the rate of cataract progression.

In addition, the PI label includes the following language regarding adverse reactions:

The most common adverse reactions (incidence $\geq 5\%$) in the clinical trials were conjunctival hyperemia, cataract, increased intraocular pressure, retinal tear, dellen (thinning of the corneal stroma), macular hole, subretinal deposits, eye inflammation, eye irritation, eye pain, and maculopathy (wrinkling on the surface of the macula).

Definitions

Biallelic Mutation: A mutation in both copies of a particular gene that affects the function of both copies.

Mutation: A change in the usual DNA sequence of a particular gene; the change can be harmful, beneficial or neutral.

Retina: Light sensitive tissue located at the back of the eye which transmits images along the optic nerve to the brain.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Voretigene neparvovec-rzyl (Luxturna)

Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

When services may be Medically Necessary when criteria are met:

HCPCS

J3398 Injection, voretigene neparvovec-rzyl, 1 billion vector genomes [Luxturna]

ICD-10 Diagnosis

H35.50-H35.54 Hereditary retinal dystrophy

When services are Investigational and Not Medically Necessary:

For the procedure code listed above when criteria are not met or for all other diagnoses not listed; or when the code describes a procedure indicated in the Position Statement section as investigational and not medically necessary.

References

Peer Reviewed Publications:

1. Ashtari M, Zhang H, Cook PA, et al. Plasticity of the human visual system after retinal gene therapy in patients with Leber's congenital amaurosis. *Sci Transl Med.* 2015; 7(296):296ra110.
2. Bainbridge JW, Mehat MS, Sundaram V, et al. Long-term effect of gene therapy on Leber's congenital amaurosis. *N Engl J Med.* 2015; 372(20):1887-1897.
3. Bainbridge JW, Smith AJ, Barker SS, et al. Effect of gene therapy on visual function in Leber's congenital amaurosis. *N Engl J Med.* 2008; 358(21):2231-2239.
4. Bennett J, Ashtari M, Wellman J, et al. AAV2 gene therapy readministration in three adults with congenital blindness. *Sci Transl Med.* 2012;4(120):120ra15.
5. Bennett J, Wellman J, Marshall KA, et al. Safety and durability of effect of contralateral-eye administration of AAV2 gene therapy in patients with childhood-onset blindness caused by RPE65 mutations: a follow-on phase 1 trial. *Lancet.* 2016; 388(10045):661-672.
6. Cideciyan AV, Aleman TS, Boye SL, et al. Human gene therapy for RPE65 isomerase deficiency activates the retinoid cycle of vision but with slow rod kinetics. *Proc Natl Acad Sci U S A.* 2008; 105(39):15112-15117.
7. Cideciyan AV, Jacobson SG, Beltran WA, et al. Human retinal gene therapy for Leber congenital amaurosis shows advancing retinal degeneration despite enduring visual improvement. *Proc Natl Acad Sci U S A.* 2013; 110(6):E517-E525.
8. Dalkara D, Goureau O, Marazova K, Sahel JA. Let there be light: gene and cell therapy for blindness. *Hum Gene Ther.* 2016; 27(2):134-147.
9. Di Iorio V, Karali M, Brunetti-Pierri R, et al. Clinical and genetic evaluation of a cohort of pediatric patients with severe inherited retinal dystrophies. *Genes (Basel).* 2017; 8(10).
10. Dias MF, Joo K, Kemp JA, et al. Molecular genetics and emerging therapies for retinitis pigmentosa: basic research and clinical perspectives. *Prog Retin Eye Res.* 2018; 63:107-131.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Voretigene neparvovec-rzyl (Luxturna)

11. Hauswirth WW, Aleman TS, Kaushal S, et al. Treatment of Leber congenital amaurosis due to RPE65 mutations by ocular subretinal injection of adeno-associated virus gene vector: short-term results of a phase I trial. *Hum Gene Ther.* 2008; 19(10):979-990.
12. Jacobson SG, Cideciyan AV, Ratnakaram R, et al. Gene therapy for Leber congenital amaurosis caused by RPE65 mutations: safety and efficacy in 15 children and adults followed up to 3 years. *Arch Ophthalmol.* 2012; 130(1):9-24.
13. Jacobson SG, Cideciyan AV, Roman AJ, et al. Improvement and decline in vision with gene therapy in childhood blindness. *N Engl J Med.* 2015; 372(20):1920-1926.
14. Kumaran N, Moore AT, Weleber RG, Michaelides M. Leber congenital amaurosis/early-onset severe retinal dystrophy: clinical features, molecular genetics and therapeutic interventions. *Br J Ophthalmol.* 2017; 101(9):1147-1154.
15. Le Meur G, Lebranchu P, Billaud F, et al. Safety and long-term efficacy of AAV4 gene therapy in patients with RPE65 Leber congenital amaurosis. *Mol Ther.* 2018 Jan 3;26(1):256-268.
16. Maguire AM, High KA, Auricchio A, et al. Age-dependent effects of RPE65 gene therapy for Leber's congenital amaurosis: a phase 1 dose-escalation trial. *Lancet.* 2009; 374(9701):1597-1605.
17. Pierce EA, Bennett J. The status of RPE65 gene therapy trials: safety and efficacy. *Cold Spring Harb Perspect Med.* 2015; 5(9):a017285.
18. Ripamonti C, Henning GB, Robbie SJ, et al. Spectral sensitivity measurements reveal partial success in restoring missing rod function with gene therapy. *J Vis.* 2015;15(15):20.
19. Russell S, Bennett J, Wellman JA, et al. Efficacy and safety of voretigene neparvovec (AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a randomised, controlled, open-label, phase 3 trial. *Lancet.* 2017; 390(10097):849-860.
20. Scholl HP, Moore AT, Koenekoop RK, et al; RET IRD 01 Study Group. Safety and proof-of-concept study of oral QLT091001 in retinitis pigmentosa due to inherited deficiencies of retinal pigment epithelial 65 protein (RPE65) or lecithin:retinol acyltransferase (LRAT). *PLoS One.* 2015; 10(12):e0143846.
21. Simonelli F, Maguire AM, Testa F, et al. Gene therapy for Leber's congenital amaurosis is safe and effective through 1.5 years after vector administration. *Mol Ther.* 2010; 18(3):643-650.
22. Testa F, Maguire AM, Rossi S, et al. Three-year follow-up after unilateral subretinal delivery of adeno-associated virus in patients with Leber congenital Amaurosis type 2. *Ophthalmology.* 2013; 120(6):1283-1291.
23. Weleber RG, Pennesi ME, Wilson DJ, et al. Results at 2 years after gene therapy for RPE65-deficient Leber congenital amaurosis and severe early-childhood-onset retinal dystrophy. *Ophthalmology.* 2016; 123(7):1606-1620.
24. Wright AF. Long-term effects of retinal gene therapy in childhood blindness. *N Engl J Med.* 2015; 372(20):1954-1955.

Government Agency, Medical Society, and Other Authoritative Publications:

1. American Academy of Ophthalmology (AAO). Clinical Statement. Recommendations on Clinical Assessment of Patients with Inherited Retinal Degenerations- 2016. For additional information visit the AAO website: <https://www.aao.org/guidelines-browse?filter=clinicalstatement> . Accessed on March 5, 2018.
2. Food and Drug Administration (FDA). FDA Advisory Committee Briefing Document: Spark Therapeutics, Inc LUXTURNA™ (voretigene neparvovec). October 27, 2017. Available at: <https://www.fda.gov/downloads/advisorycommittees/committeesmeetingmaterials/bloodvaccinesandotherbiologics/cellulartissueandgenetherapiesadvisorycommittee/ucm579300.pdf>, Accessed on February 12, 2019.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member's contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Voretigene neparvovec-rzyl (Luxturna)

3. Luxturna [Product Information], Philadelphia, PA. Sparks Therapeutics, Inc; Available at: <https://www.fda.gov/downloads/BiologicsBloodVaccines/CellularGeneTherapyProducts/ApprovedProducts/UCM589541.pdf>. Accessed on February 12, 2019.
4. Stone EM, Aldave AJ, Drack AV, et al. Recommendations for genetic testing of inherited eye diseases: report of the American Academy of Ophthalmology task force on genetic testing. *Ophthalmology*. 2012; 119(11):2408-2410.

Websites for Additional Information

1. American Association for Pediatric Ophthalmology and Strabismus (AAPOS). Leber’s Congenital Amaurosis. Updated on January 2018. Available at: <https://www.aapos.org/terms/conditions/66>. Accessed on February 12, 2019.
2. National Institute for Health: National Eye Institute (NEI). Gene Therapy for Leber Congenital Amaurosis. Available at: <https://nei.nih.gov/lca>. Accessed on February 12, 2019.
3. National Institute of Health: U.S. National Library of Medicine. RPE65 gene. Reviewed August 2010. Available at: <https://ghr.nlm.nih.gov/gene/RPE65>. Accessed on February 12, 2019.
4. National Institute of Health: U.S. National Library of Medicine. What is gene therapy? Reviewed February 12, 2019. Available at: <https://ghr.nlm.nih.gov/primer/therapy/genetherapy>. Accessed on February 12, 2019.

Index

SPK-RPE65

The use of specific product names is illustrative only. It is not intended to be a recommendation of one product over another, and is not intended to represent a complete listing of all products available.

Document History

Status	Date	Action
Reviewed	03/21/2019	Medical Policy & Technology Assessment Committee (MPTAC) review. Revised title from [™] to [®] symbol. Updated References.
	12/27/2018	Updated Coding section with 01/01/2019 HCPCS changes; added J3398, removed C9032 deleted 12/31/2018.
	06/28/2018	Updated Coding section with 07/01/2018 HCPCS changes; added C9032.
New	03/22/2018	Updated Rationale to include information regarding pediatric population. MPTAC review. Initial document development.

Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The member’s contract benefits in effect on the date that services are rendered must be used. Medical Policy, which addresses medical efficacy, should be considered before utilizing medical opinion in adjudication. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.